

WILLOWOOD CLORAN DF

Same active as *FIRSTRATE*®



OVERVIEW

Willowood Cloran DF is a highly effective pre emergence and post emergence herbicide with residual control on the most troublesome broadleaf weeds including glyphosate-resistant species.

KEY CROPS

Soybeans

PACKAGING

- 10 oz package (x8)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Cloransulam-Methyl (84% Active Ingredient per Gallon)

RATES

0.3 oz/A - 0.75 oz/A
(See label for rates by crop)

WEED SPECTRUM

Marestail • Waterhemp • Lambsquarters • Ragweed • Pigweed

APPLICATION TIMING

Preplant Incorporated • Preplant • Burndown • Pre Emergence • Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • ALS INHIBITORS
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Wheat: 4 months • Corn: 9 months
Adjuvant Recommendation	<ul style="list-style-type: none"> • NIS at .25% V/V or COC/MSO at 1.2% V/V • UAN at 2.5% V/V or AMS at 2lb/A
Rainfastness	<ul style="list-style-type: none"> • 2 hours
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Minimum of 3GPA
Application Tips	<ul style="list-style-type: none"> • Good at controlling Giant Ragweed in Conventional or GT Soybeans post emergence. • Refer to the annual maximum use rate of Willowood Cloran DF when using any other product containing Cloransulam as a pre emergence application.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.



WILLOWOOD CLORAN DF

Same active as *FIRSTRATE*®

MIXING PROCEDURE

Mix Willowood Cloran DF as follows:

- Fill the spray tank to 1/4 to 1/3 of the total spray volume required with water or liquid fertilizer.
- Start agitation.
- Add the required amount of Willowood Cloran DF for the acreage being treated directly to the spray tank while agitating and allow time to disperse. If liquid fertilizer is being used as the spray carrier rather than water, pre-mix Willowood Cloran DF as described above before adding to the spray tank.
- After adding Willowood Cloran DF, add different formulation types in the following order: other formulation(s) packaged in water soluble packets; any compatibility agent, if required; dry flowables; wettable powders; aqueous suspension, flowable and liquids.
- Maintain agitation and fill spray tank to 3/4 of total spray volume and add:
 - Emulsifiable concentrates; solutions; adjuvants. Allow time for complete mixing and dispersion after each addition.
- Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application.

CROP STAGING

- Preplant/Pre Emergence: apply up to two weeks prior to planting or pre emergence for additional burndown and residual control.
- Post Emergence: may be applied any time prior to the R2 (full flower) growth stage of soybeans.

FIELD NOTES
